

What Future Expects Humanity After the Demographic Transition Time?

L.Ya.Kobelev, L.L.Nugaeva
 Department of Physics, Urals State University
 Lenina Ave., 51, Ekaterinburg 620083, Russia
 E-mail: leonid.kobelev@usu.ru

The variant of phenomenological theory of humankind future existence after time of demographic transition based on treating the time of demographic transition as a point of phase transition and taking into account an appearing of the new phase of mankind is proposed. The theory based on physical phenomenological theories of phase transitions and classical equations for system predatory-preys for two phases of mankind, take into account assumption about a multifractal nature of the set of number of people in temporal axis and contains control parameters. The theory includes scenario of destroying of existent now human population by new phase of humanity and scenario of old and new phases co-existence. In particular cases when the new phase of mankind is absent the equations of theory may be formulated as equations of Kapitza, Foerster, Hoerner, Kobelev and Nugaeva, Johansen and Sornette phenomenological theories of growth of mankind.

01.30.Tt, 05.45, 64.60.A; 00.89.98.02.90.+p.

I. INTRODUCTION

The problem of mankind population growth is one of the global problems of the mankind existence and its prosperity in the future. Will the demographic explosion existing now at the mankind population of the whole world stops and what will be after its stopping? Almost all the phenomenological theories of future of mankind (see [1], [2], [3], [4], [5]) include the time of demographic transition t_0 . Before this time the growth of mankind population is very large (hyperbolic or exponential growth), after this time the population growth may practically absent [3] or may be regulated by driving parameters [4] that govern the development of mankind (such parameters are present in non-linear open systems (see [6])). Thus the time point t_0 (the time of demographic transition) is very essential for the future of mankind. What is biological and physical senses of the demographic transition time? The mankind as the whole is a very large system and its behavior can be described by methods of nonlinear dynamic, probability theory or statistical theory of open system using the methods of fractal geometry ([7], [8], or power laws [5]). Now put the question: how the demographical transition time may be treated from these points of view? There are many different ways to answer on this question. In this paper we consider the demographic transition time as a point of a phase transition, i.e. the point of transition of mankind in a new phase (in physiological, psychological, behaviour and so on senses). What this phase will be if it may exist? Will be the population of men of the new phase friendly to the old phase of mankind or will they fight with men of the old phase and force out them from their ecological niche? What scenario may exist and be admissible in that case? It seems that if the driving parameters that determine growth of the new phase (thou it is difficult

now to select the main of them, it is the task for many research groups) will be the same as now the new phase may diminish the old phase of mankind very soon. We would remind that all phenomenological theories give for the time t_0 the time in first part of twenty one century. If the treating of time t_0 as a point of phase transition is correct and has relation to existence of humanity, the embryos of the new phase must appear (as it well known from theory of phase transitions in physics) before the time of demographic transition, i.e. in very near time. What role they will play in the life of mankind society and what it will be like? Will the new phase consist of international groups of islam terrorist and the fight and wars spread to the whole world? Or it will international groups of corrupted officials that for own prosperity govern the whole world to its death? May be the groups of sexual minorities will became the main part of mankind and diminish it? Will the new phase characteristics lay in psychological (i.e. behaviour) domain or it lay in the physiological domain (may be it is incurable genetics diseases or mental disorders reasoned by increasing radiation in our world)? Are the Russian revolution in 1917 year or the Germany fascist state in 1933 year are the examples of appearing of such phases? May be the victory of progressive mankind in the World War II and collapse of socialist system in Soviet Union are results only the fact that the demographic time transition is not come? It seems, the search of answers on these questions is the one of main tasks of governments and research demographical gropes of developed countries in near time.

We consider in this paper only one way of mathematical analysis and description of two phases of mankind (the old phase and the new phase) and describe time behaviour of these phases by system of differential equations that are Volterra-Lotka equations for system of

predatory-prey.

II. EQUATIONS FOR DESCRIBING TWO PHASES MANKIND TIME BEHAVIOUR

There are two simplest ways (physical and biological) for describing time dependence of two phases (the old and the new) of mankind :

1. In physics the phenomenological theory of phase transitions was developed by Landau (see [9]). Theory of phase transitions as appearing of order parameters developed by Ginzburg-Landau [10]. These theory may be used for describing time behaviour of two phases. From physical point of view the transition from the old phase of mankind to the new phase may: a) take very short time (as transition of water to ice); b) take long time and has intermediate state when two phases of mankind will co-existent (as in the transition to superconductive state for superconductors of second order); c) be characterized by stretched in time axes phase transition as in some ferroelectrics. The differences of these cases determined by characteristics of the small embryos gropes of new phase and their interaction with the old phase of mankind. If interaction between men of old and new phases will attract the man of old phase to transition to the new phase (we will call such interaction as "negative"), the transition time will be small and co-existent of two phases of mankind for the long time period is impossible. Probably the evidences of such transitions are the above examples of Russian revolution of 1917 year and fascist Germany state of 1933 year. If result of interaction of men of the old phase with gropes of men of the new phase will neutral, i.e. men of old phase will not be attracted by embryos of new phase ("positive" interaction), amount of new the phase will increase and co-existence of new and old phases is possible. The time of coexistence determined by govern parameters and selected mathematical models.

2. The biological point of view on behavior of mankind allows to use many mathematical models for describing of time behaviour of old and new mankind phases. In this paper we see only simple predatory-prey models where the old phase of mankind will be prey and the new phase will be predatory.

III. EQUATIONS OF WOLTERRA-LOTKA FOR OLD AND NEW PHASES MANKIND POPULATION

Let us designate the old phase of mankind by $N_1(t)$ and the new phase of mankind by $N_2(t)$ where t is current time. Equations of Volterra-Lotka type is well researched. It is well considered the time behaviour of the

predatory and the prey populations (including time behaviour populations near the point of demographic transition t_0 where fluctuations play important role). So we only write these equations and mark the main consequences of mathematical theories. Let for time dt the old phase changes are $cN_1^\nu dt - \gamma N_1 N_2 dt$ (the second member is the result of diminishing character of influences (interactions) between the old and the new phases) and new phase changes are $-\delta N_2 dt + b N_1 N_2 dt$. The equations for $N_1(t)$ and $N_2(t)$ may be written in this case as

$$\frac{\partial}{\partial t} N_1(t) = cN_1^\nu(t) - \gamma N_1(t)N_2(t) \quad (1)$$

$$\frac{\partial}{\partial t} N_2(t) = -\delta N_2(t) + b N_1(t)N_2(t) \quad (2)$$

where c, b, δ, γ are constant and defined by selections of considered models. These equations are Volterra equations for $\nu = 1$ and are Lotka equations for $\nu = 0$. The humanity phases N_1 and N_2 in (1),(2) have links and determine one another.

In the case when populations of the old phase and the new phase are multifractal sets on the time axes (see [4]) equations (1) and (2) read

$$D_{0,t}^\beta N_1(t) = cN_1^\nu(t) - \gamma N_1(t)N_2(t) \quad (3)$$

$$D_{0,t}^\mu N_2(t) = -\delta N_2(t) + b N_1(t)N_2(t) \quad (4)$$

where $D_{0,t}^\beta$ and $D_{0,t}^\mu$ are Riemann-Liouville fractional derivatives or generalized Riemann-Liouville derivatives (see [4]) and β and μ are fractional numbers or functions of govern parameters.

The equations (1) and (2) may be considered as the base equations of rude biological models for describing future population of mankind if there are exist two phases of mankind and sets of N_1 and N_2 are sets of population of old and new phases . As well known the equations of such type (type of Volterra-Lotka equations) may be generalized and wrote in more general form (form of Kolmogorov-Fokker-Plank equations for probability density $w(N_1, N_2, t)$ to find in moment t the populations N_1 and N_2 in the considered system [12], [13], or may be considered the case when influences N_2 on behaviors of N_1 included only at the bound time intervals and so on). These generalized equations are wide spread and give many known models have used not only in biology ([14]), but in economics ([15]), non linear physics, theoretical chemistry and so on. In the next part of paper we write more detailed description of results of interactions between two mankind phases N_1 and N_2 based on well known researches of equations of such type. For case when in above equations there are only one phase N_1 , and $\nu = 0$ and coefficient c is function of time and t_0 ($c = c(t, t_0)$) it is not difficult to chose $c(t, t_0)$ in the simple forms and as special cases receive the base equations of work [3], [4], [5].

IV. HOW WILL INTERACT OLD AND NEW MANKIND PHASES?

The mathematical research of solutions characteristics of equations (1)-(2) or more general equations taking into account the fluctuations is well worked out (see for example [13]. As was pointed out these equations may be written for mean values of N_1 and N_2 or in the form of govern equations for probability $w(N_1, N_2, t)$ to find the system in the moment t with values N_1 and N_2 , or in the form of Kolmogorov- Fokker-Plank equations that are consequences of governor equations. The influences of noises and fluctuations(depending at c, b, δ, γ) on the behaviour of system may be taken into account too. More complicated equations may be written for case when the phase N_2 may part of her time have "a rest time" and in that rest time do not interact with the phase N_1 . So we omit the process of receiving of solutions for different systems of equations describing interaction between old-new phases and only describe main results:

1. Possibility of co-existence between old and new phases with periodical changing of value of N_1 and N_2 ;
2. Possibility for diminishing of one phase by other phase that depends at the initial values;
3. Possibility for calculation of mean values of N_1 and N_2 in one of system states;
4. Possibility for calculation of time existence t_{ex} for old or new phases if take into account by including in the main equations (or changing the main equation by its Fokker-Plank probability analogies equations) external or internal noises (additive or multiplicative);
5. Possibility of description of fluctuation characteristics of kinetic variables near bifurcation points;
6. Possibility of researching of noises role in losing stability of stationary state or in different self-excited oscillations states;
7. Possibility of determination of conditions for stochastic regimes appearing in system;
8. Possibility of governing by processes of fighting or co-existence between old and new phases;
9. Possibility of using more complicated models and equations including fractal characteristics of mankind population distribution on the time axis.
10. Possibility of selection of different variants for govern parameters based on experimental sociological dates and so on.

Thus the predatory-prey models and its equations give rich possibilities for researching of the behavior of two phases of mankind. Has this assumption about appearing of two mankind phases relation to problem of future propagation of mankind? The future propagation of mankind will gives answer on this question, but if it assumption is correct it is time now to find the new phase (or phases) and determine (or make attempts to determine) is it one or other phases useful for future of mankind or not useful.

V. CONCLUSIONS

As one of advantages to use the equations (1)-(2) for describing demographic problems (with some of them the mankind already has confronted now and may be these problems are consequences of appearing the embryos of new phase of mankind) we shall stress an opportunity of insert and account in the theory many factors (such as, incurable illnesses, natural cataclysmic etc.) defining future of mankind as a result of influences the control parameters included in the govern parameters of predatory-prey models and fractional dimensions .

There are three main scenario of models based on equations type of (1)-(2):

- a) Two phases of mankind co-existent during long time with periodically changes its values and with mean values $N_1 = \delta b^{-1}$ and $N_2 = c\gamma^{-1}$ conserves in bound limits if population fluctuations may be omitted;
- b) Mankind population lost its stability as the result of fluctuation influences if the equations of Lotta type used;
- c)The old mankind phase has bound time of existence and this time may by calculated if parameters of theory are defined .

The main purpose of this paper was to analyze possibility of introducing of two mankind's population phases after demographic transition time. Driving parameters included in the considered models (equations of Landau-Ginzburg time theory or equations of predatory-prey type) of the phenomenological theories of the mankind population in the frame of physical Landau or Ginzburg-Landau models or biological predatory-prey models. The predatory-prey type models was chosen asan example. The change of number of mankind (described in the framework of phenomenological theories of the population) can be adjusted by such choice of control parameters with concrete contents of dependencies of them at which the mankind will has stability time and population will grow so slowly, that overpopulation and the problems connected with fighting between two mankind phases will do not arise in the foreseen future. Stress now as was done in [5] that after demographic transition time also the type of economical development of humankind may be changed .

-
- [1] Foerster, Von H. et al., Science, v.132, p.1291 (1960)
 - [2] Hoerner, von S.J., British Interplanetary Society, v.28, 691, (1975)
 - [3] Kapitza S.P., Uspehi Fizicheskikh Nauk (Russia), 1996, vol.166, No.1, pp.63-79; Kapitza S.P., *How Many People Lived, Live and are to Live in the World. An essay on the theory of growth of humankind*, Moscow, Inst. Phys. Problem RAS, 1999,238p.(in Russian)

- [4] Kobelev L.Ya., Nugaeva L.L. Will Humanity population Stabilized in Future? arXiv/physics:0003035
- [5] Johansen A., Sornette D. The End of growth Era? arXiv:cond-mat/0002075
- [6] Klimontovich Yu.L., *Statistical Theory of Open Systems. Vol.1*, Moscow, Yanus, 1995, 686p. (in Russian); Kluwer Academic Publishers, Dordrecht, 1995; Klimontovich Yu.L., *Statistical Physics of Open systems. Vol.2*, Moscow, Yanus, 1999, 450p. (in Russian).
- [7] Mandelbrot B., *Fractal Geometry of Nature*, W.H.Freeman, San Francisco, 1982
- [8] Kobelev L.Ya, *Fractal Theory of Time and Space*, Ekaterinburg, Konros, 1999, 136p. (in Russian); Kobelev L.Ya., What Dimensions Do the Time and Space Have: Integer or Fractional? arXiv:physics/0001035;
- [9] L.D.Landau, I.M.Livshiz, *Statistical Physics, Moscow, Nauka, 1964 (in Russian)*
- [10] V.L.Ginzburg, L.D.Landau, GETP, 1948, v.20,P.1064 (in Russian)
- [11] Samko S.G, Kilbas A.A., Marichev I.I., *Integrals and Derivatives of the Fractional Order and Their Applications*, (Gordon and Breach, New York, 1993).
- [12] Alekseev B.B.,Kostin I.K,Stability of simple biogeochinoses to External Stochastic Influences,in "Biological Systems in Agriculture and Forestry",-Moskwa, Nauka,1974, p.105-112 (in Russian)
- [13] Vasiliev,V.A., Romanovski Yu.M. Yaxno V.G. Avtovave processes, Moscow, Nauka,1987,p.240 (in Russian)
- [14] Murray J.D. *Mathematical Biology*, Springer-Verlag, Berlin New York Heidelberg, 1989
- [15] Lorenz *Nonlinear Dynamical Economics and Chaotic Motion*, Springer-Verlag, 1993.